

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

others seemed larger and grayer than the White-throats. This is a rare bird in our area.—NATHAN F. LEOPOLD, *4754 Greenwood Ave., Chicago, Ill.*

Rare Birds in the Indiana Sand Dunes.—Picoides arcticus. ARCTIC THREE-TOED WOODPECKER.—A fine male of this species was taken one mile west of the Dune Park station in Porter County, Indiana, October 3, 1920. Another, also a male, was taken a mile and a half east of Dune Park on October 24, 1920. These specimens constitute the second and third Indiana records and were taken within four miles of where the first specimen was taken March 11, 1917 ('Auk,' Oct. 1917, pp. 487). The specimen taken October 24 appeared rather shy and wild, flying nervously from tree to tree uttering a loud and rather startling note, a rapidly repeated 'teck-teck-teck.' The other was extremely tame, industriously digging grubs out of a small dead scrub pine from which he had the bark almost stripped. It seems remarkable that these northern birds should appear so far south before cold weather had set in.

Zonotrichia querula. HARRIS'S SPARROW.—An immature female of this species was taken at Millers, Lake County, Indiana, on October 3, 1920.

Spizella pallida. CLAY-COLORED SPARROW.—A female was taken at Dune Park, Porter County, Indiana, on May 25, 1919. These records are the second for Indiana in each case to the best of my knowledge. Careful examination of the large migrating flocks of sparrows in the northwestern part of the State should reveal a few Harris's Sparrows as it does all along the eastern border of their range. A specimen was seen about fifty miles north of here, in Jackson Park, Chicago, September 30, a circumstance that caused me to look for them in the Dunes.

The skins of the specimens taken are in the Harris Extension Collection.—H. L. STODDARD, *N. W. Harris Public School Extension of Field Museum, Chicago, Ill.*

Additions to the Birds of Lake County, Minnesota.—A visit to Lake County during the latter part of the summer of 1920 resulted in the following additions to my list, published in 'The Auk,' October, 1920, p. 541:

Colymbus auritus (Linn.). HORNED GREBE.—August 6 three Horned Grebes were observed at close range near the south shore of Lake Bald Eagle.

Pisobia maculata (Vieill.). PECTORAL SANDPIPER.—August 25 four specimens were shot and about two dozen others observed on a mud-flat in the Isabelle River some distance below Rice Lake.

Pelidna alpina sakhalina (Vieill.). RED-BACKED SANDPIPER.—August 25 one of these birds, in the company of four Yellow-legs, was shot along the Isabelle River about a mile below Rice Lake.

Ereunetes pusillus (Linn.). SEMIPALMATED SANDPIPER.—August 20 one was shot out of a flock of four on the east shore of Lake Isabelle. On